arXiv:1212.5712v1 [physics.bio-ph] 22 Dec 2012

Efficient fold-change detection based on protein-protein interactions

Wouter Buijsman and Michael Sheinman
Department of Physics and Astronomy, VU University, 1081 HV Amsterdam, The Netherlands

(Dated: December 27, 2012)

Various biological sensory systems exhibit a response to the relative change of the stimulus, often reffered to as fold-change detection. Here, we present a mechanism consisting of two interacting proteins, able to detect a fold-change effectively. This mechanism, in contrast to other proposed mechanisms, does not consume chemical energy and is not subject to transcriptional and translational noise. We show by analytical and numerical calculations that the mechanism can have a fast, precise and efficient response for parameters that are relevant to eukaryotic cells.

PACS numbers:

I. INTRODUCTION

A fold-change detecting (FCD) mechanism with input u and output y is characterized by the property

$$y \propto \frac{\Delta u}{u}, \tag{1}$$

indicating that the output signal depends only on a relative change in the input signal rather than the absolute value [1]. FCD fits various biologial sensory systems, such as vision and hearing. Recent studies show that FCD mechanisms are present in *E. coli* [2] and *Xenopus laevis* embryos [3].

In the last few years, it was demonstrated that the incoherent feedforward loop [4, 5] and the two-state protein [6] can provide FCD. As a consequence of the involvement of continues protein production and degradation, they consume chemical energy, and are subject to transcriptional and translational noise [7]. Besides, transcription and translation set a lower limit for the response time, which can be quite high, especially in mammalian cells (∼ 20min) [8–10].

Here, we describe a FCD mechanism, based purely on protein-protein and protein-ligand interactions. The mechanism it is not subject to transcriptional and translational noise and does not consume chemical energy since it does not involve protein production and degradation. The characteristic timescales are set by the rates of protein interactions, which have a much broader range than the transcription and translation rates [11–14].

This paper contains three sections. Section II describes the mechanism and shows formally that the mechanism can provide FCD. In section III, the action of the mechanism is demonstrated by numerical simulation. In this section, it is shown that the mechanism can be preciese, robust and efficient for biologically relevant parameters. Section IV provides a discussion and outlook of the results.

II. THE MECHANISM

The mechanism we suggest uses two proteins, named X and Y. The system responds with the output Y_S to the concentration of the input ligand, named S, where Y_S is the complex of Y and S. The mechanism involves three reactions, as presented schematically in Fig 1. In the first reaction, S and X bind, forming the complex X_S. The binding and unbinding rates are given by $k^{\text{on}}_{X_S}$ and $k^{\text{off}}_{X_S}$, respectively:



FIG. 1: A schematic representation of the FCD mechanism as proposed in this paper. The mechanism has input S and output Y_S. For clarity, Y is drawn two times. The copies are grouped by a box.

$$S + X \underset{k^{\text{off}}_{X_S}}{\overset{k^{\text{on}}_{X_S}}{\rightleftarrows}} X_S. \tag{2}$$

In the second reaction, S and Y bind, forming the output complex Y_S with binding rate $k^{\text{on}}_{Y_S}$ and unbinding rate $k^{\text{off}}_{Y_S}$:

$$S + Y \underset{k^{\text{off}}_{Y_S}}{\overset{k^{\text{on}}_{Y_S}}{\rightleftarrows}} Y_S. \tag{3}$$

In the third reaction, X_S and Y bind, forming the complex X_SY with binding and unbinding rates given by $k^{\mathrm{on}}_{X_SY}$ and $k^{\mathrm{off}}_{X_SY}$, respectively:

$$X_S + Y \underset{k^{\mathrm{off}}_{X_SY}}{\overset{k^{\mathrm{on}}_{X_SY}}{\rightleftarrows}} X_SY. \tag{4}$$

Fig. 1 shows a schematic representation of the mechanism.

A. The mechanism satisfies FCD conditions

In general, a mechanism based on input S, internal variable X, and output Y_S, described by functions

$$\dot{X}_S = f(X_S, Y_S, S), \tag{5}$$

$$\dot{Y}_S = g(X_S, Y_S, S). \tag{6}$$

provides FCD if [5]:

$$f(p\,X_S, Y_S, p\,S) = p\,f(X_S, Y_S, S), \tag{7}$$

$$g(p\,X_S, Y, p\,S) = g(X_S, Y_S, S), \tag{8}$$

for any $p > 0$. Below, we show that the suggested mechanism can provide FCD. This is accomplished by showing that conditions (7) and (8) can be satisfied.

1. *First condition,* (7)

The first condition, (7), requires the ratio S/X_S for an input S, which increases by a factor ϕ,

$$S(t) = \begin{cases} S_0 & \text{if } t < t_0 \\ \phi\,S_0 & \text{if } t \geq t_0 \end{cases}, \tag{9}$$

to be independent of the initial input concentration S_0. When concentration Y is small compared to concentrations S and X, the major reaction that involves X_S is the binding of X and S given by reaction (2). The mean-field concentration X_S as a function of time is described by the Michaelis-Menten equation [15], which for $k^{\mathrm{on}}_{X_S}X \ll k^{\mathrm{off}}_{X_S}$ can be approximated by

$$\dot{X}_S = k^{\mathrm{on}}_{X_S} X S - k^{\mathrm{off}}_{X_S} X_S. \tag{10}$$

By comparison with condition (7), it follows that, in this case, the first condition is satisfied.

2. *Second condition,* (8)

The second condition, (8), requires the output Y_S to be proportional to the ratio S/X_S. When reactions (3) and (4) equilibrate fast compared to reaction (2), a continues equilibrium for reactions involving Y can be assumed. If Y has a strong affinity to X_S, it follows from the Michaelis-Menten equation that the equilibrium value of Y_S is approximated by

$$Y_S = \frac{S/k_{Y_S}}{X_S/k_{X_SY}} Y, \tag{11}$$

where $k_{X_SY} = k^{\mathrm{off}}_{X_SY}/k^{\mathrm{on}}_{X_SY}$ and $k_{Y_S} = k^{\mathrm{off}}_{Y_S}/k^{\mathrm{on}}_{Y_S}$. By comparison with condition (8), it follows that, in this case, the second condition is also satisfied.



FIG. 2: Mean-field concentrations for $S = 10nM$, $X = \mu M$, $Y = 10nM$, $k^{\mathrm{on}}_{X_S} = 10^7 M^{-1}s^{-1}$, $k^{\mathrm{off}}_{X_S} = 10^{-1}s^{-1}$, $k^{\mathrm{on}}_{Y_S} = 10^9 M^{-1}s^{-1}$, $k^{\mathrm{off}}_{Y_S} = 10s^{-1}$, $k^{\mathrm{on}}_{X_SY} = 10^9 M^{-1}s^{-1}$, and $k^{\mathrm{off}}_{X_SY} = 10s^{-1}$ as a function of time. Here, $1nM$ corresponds to 1 protein for prokaryotes and 10^3 proteins for eukaryotes. The input S fold-changes at $t = 1.5s$ and $t = 4s$ by a factor $\phi = 3$.

B. Mean-field FCD

In sum, it was shown that the mechanism provides FCD if

a. concentrations S and X are large compared to Y;

b. reaction rate $k^{\mathrm{on}}_{X_S}X$ is small compared to $k^{\mathrm{off}}_{X_S}$;



FIG. 3: Mean-field solution and random walk simulation of protein concentrations for $S = 10nM$, $X = \mu M$, $Y = 10nM$, $k^{\text{on}}_{X_S} = 10^7 M^{-1}s^{-1}$, $k^{\text{off}}_{X_S} = 0.1s^{-1}$, $k^{\text{on}}_{Y_S} = 10^9 M^{-1}s^{-1}$, $k^{\text{off}}_{Y_S} = 10s^{-1}$, $k^{\text{on}}_{X_SY} = 10^9 M^{-1}s^{-1}$, and $k^{\text{off}}_{X_SY} = 10s^{-1}$. The input S is presented on the upper plots. It fold-changes at $t = 2, 4.5, 7s$ and $t = 9.5s$ by a factor $\phi = 3$. On the bottom plots mean-field solution is represented by the thick gray lines while thin black lines represent random walk simulations. For eukaryotes, the two lines mostly overlap. Here, $1nM$ corresponds to 1 protein for prokaryotes and 10^3 proteins for eukaryotes.

c. protein Y has a strong affinity to X_S compared to S;

d. reactions (3) and (4) equilibrate fast compared to reaction (2).

When the conditions are satisfied, the FCD transient response is characterized by two timescales,

$$\tau_1 = \frac{Y_s}{k^{\text{on}}_{Y_S} Y S} \text{ and } \tau_2 = \frac{X_S}{k^{\text{on}}_{X_S} X S}, \qquad (12)$$

corresponding to the raising and the decay of the response, respectively.

Fig. 2 shows an example of mean field values X, X_S, Y, Y_S, and X_SY as a function of time for an input given by (9) with $\phi = 3$ and $t_0 = 1.5s$ and $t_0 = 4s$. One can see that the response, Y_S, does not depend on the absolute value of the input, S. This shows that themechanism provides FCD for this set of parameters. The parameter values and the robustness to noise are discussed in Sec. III.

III. EXAMPLES AND BIOLOGICAL RELEVANCE

Affinity constants of protein-protein and ligand-receptor interactions span, over six orders of magnitude, $10^{-12} - 10^{-6}M$, a result of the large range of the association and dissociation rates which are typically in the $10^3 - 10^{10} M^{-1}s^{-1}$ and $10^{-4} - 10^4 s^{-1}$ range, respectively [11–14]. Protein concentrations typically vary in the range $1 - 10^3 nM$, corresponding to $10^3 - 10^6$ proteins for eukaryotes and $1 - 10^3$ proteins for prokatyotes [16]. Below, we show that such a large range of biochemical parameters enables fulfilment of the FCD conditions, still leaving a lot of freedom. This freedom, in principle, can be used, for instance, to tune the timecale of the detection response.

In order to demonstrate the relevancy of the mechanism to biological systems, we analyzed equations (2, 3, 4) using mean-field and numerical random walk calculations [17, 18]. Fig. 3 shows the result for a certain set of parameters for prokaryotes and eukaryotes, which differ by the number of proteins present. This parameter set is used also in Fig. 2. The mean-field response peaks of the system are approximately transient and identical, which shows that the mechanism can provide FCD. Note that the last peak is, in this case, only approximately transient since condition (b) is no longer satisfied. For prokaryotic cells, the protein number is too low for an effective FCD on the level of a single cell. For eukaryotes, the output signal is nearly unaffected by the finite number noise. In this example, the timescales of the initial response is less than a tenth of a second; the decay timescale is approximately $0.25s$.

IV. DISCUSSION AND OUTLOOK

In this paper we propose a new FCD mechanism, based purely on protein-protein and ligand-protein interactions. The mechanism, in contrast to previously proposed mechanisms [4, 6], does not require continues trancription, translation and degradation of proteins. The mechanism avoids transcriptional bursts and other noise sources due to small number bottleneck processes [7], and benefits from the typical large numbers of proteins in eukariotic cells. The initial response time, in principle, is limited only by diffusion of proteins [12], and, therefore, can be smaller than a second for a biologically relevant set of parameters, as shown in Fig. 3.

For eukaryotes, the noise due to finite number statistics does not play a significant role. However, for prokaryotes, the typical protein number is not sufficient for a precise FCD on a single cell level. Previously proposed mechanisms are relatively consuming, since the number of proteins has to be large and their lifetime has to be short in order to reduce a detection noise and make the responce quick. From this perspective, the mechanism as proposed here is advantageous, since it is acting in detailed balance [19].

We showed that the mechanism is characterized by two timescales, τ_1 and τ_2 given in Eq. (12). τ_1 characterizes the raise of the response while τ_2 characterizes its decay. The value of τ_1, in principle, is limited only by diffusion of proteins [12]. Thus, as shown in the example, presented in Fig. 3, it can be smaller than a second for a biologically relevant set of parameters.

We expect that the proposed mechanism is only a single example of a new, large class of biochemical fold-change detectors, based purely on ligand-protein and protein-protein interactions.

[1] J. Ferrell, Mol. Cell **36**, 724 (2009).

[2] L. Goentoro and M. Kirschner, Mol. Cell **36**, 872 (2009).

[3] M. Lazova, T. Ahmed, D. Bellomo, R. Stocker, and T. Shimizu, Proc. Nat. Acad. Sci. **108**, 13870 (2011).

[4] L. Goentoro, O. Shoval, M. Kirschner, and U. Alon, Mol. Cell **36**, 894 (2009).

[5] O. Shoval, L. G. Y. Hart, A. Mayo, E. Sontag, and U. Alon, Proc. Nat. Acad. Sci. **107**, 15995 (2010).

[6] T. Marquez-Lago and A. Leier, BMC Systems Biology **5**, 22 (2011).

[7] J. Raser and E. O'Shea, Science **309**, 2010 (2005).

[8] H. Schröder, U. Friese, M. Bachmann, T. Zaubitzer, and W. Müller, European journal of biochemistry/FEBS **181**, 149 (1989).

[9] H. Kimura, K. Sugaya, and P. Cook, J. Cell Biol. **159**, 777 (2002).

[10] U. Alon, *An Introduction to Systems Biology: Design Principles of Biological Circuits* (Chapman and Hall/CRC, 2006).

[11] P. Bongrand, Reports on Progress in Physics **62**, 921 (1999).

[12] E. Levy and J. Pereira-Leal, Current Opinion in Structural Biology **18**, 349 (2008).

[13] R. Alsallaq and H.-X. Zhou, Proteins: Structure, Function, and Bioinformatics **71**, 320 (2008), ISSN 1097.

[14] G. Schreiber, G. Haran, and H. Zhou, ChemInform **40**, (2009).

[15] P. Nelson, *Biological Physics: Energy, Information, Life* (W.H. Freeman and company, 2004).

[16] R. Milo, P. Jorgensen, U. Moran, G.Weber, and M. Springer, Nucl. Acid. Res. **38**, D750 (2010).

[17] J. Norris, *Markov Chains (Cambridge Series in Statistical and Probabilistic Mathematics)* (Cambridge University Press, 1998).

[18] P. Doyle and J. Snell, *Random walks and electric networks* (Mathematical Association of America, 1984).

[19] G. Lan, P. Sartori, S. Neumann, V. Sourjik, and Y. Tu, Nature Phys. **advance online publication**, (2012).